증례보고: 전두피관술 후 시행한 알렉산드라이트 제모레이저에 의한 화상

홍준식¹ · 이동락² · 모영웅^{1,3} · 강인호¹ · 신혜경¹ · 이준호¹ · 정규용¹

¹동국대학교 의과대학 경주병원 성형외과학교실, ²인제대학교 의과대학 일산백병원 성형외과학교실, ³동국대학교 대학원 성형외과학교실

A Burn after Alexandrite[®] Laser Hair Removal on a Forehead Flap: A Case Report

Joon Shik Hong, M.D.¹, Dong Lark Lee, M.D., Ph.D.², Young Woong Mo, M.D.^{1,3}, Inho Kang, M.D.¹,
Hea Kyeong Shin, M.D., Ph.D.¹, Joon Ho Lee, M.D.¹ and Gyu Yong Jung, M.D., Ph.D.¹

¹Department of Plastic and Reconstructive Surgery, Dongguk University Gyeongju Hospital, Dongguk University College of Medicine, Gyeongju, ²Department of Plastic Surgery, Inje University Ilsan Paik Hospital, Inje University College of Medicine, Goyang, ³Department of Plastic and Reconstructive Surgery, Dongguk University Graduate School of Medicine, Seoul, Korea

The nose is a complex three-dimensional structure and represents a major aesthetic focus of the face. As a gold standard for nasal soft tissue reconstruction, the 'forehead flap' provides reconstructive surgeons a robust pedicle and large amount of tissue to reconstruct almost any defect. However, during this process, some hair can be unintentionally introduced to the nose. Accordingly, laser hair removal is sometimes needed, but blood circulation and flap survival should be carefully monitored. Despite careful evaluation, a third-degree burn occurred in our patient that required eight weeks to heal. Here, we report on a burn resulting from epilation conducted 2 weeks after forehead flap for nasal reconstruction. (*J Korean Burn Soc* 2021;24:14-17)

Key Words: Alexandrite laser, Nose deformities, Burns, Hair removal

INTRODUCTION

The Alexandrite LASER (Light Amplification by Stimulated Emission of Radiation) is one of the best options for removing unwanted hair and provides a cosmetic effect relatively safely. This laser uses an alexandrite crystal to emit energy at a wavelength of 755 nm. Although clinically used for hair, tattoo, and pigmented lesion removal, the laser is most commonly used for hair removal [1]. However, when this equipment is used, the skin is inevitably burned and the wound healing process then takes several weeks dependent on blood circulation. In

other words, when the Alexandrite LASER is used on inset flaps with poor blood circulation burns would be more likely and treatment time would be extended. In this case, we report a burn that occurred during hair removal from a forehead flap required after the removal of a huge nasal congenital nevus.

CASE REPORT

A 65-year-old female patient visited our hospital with a large, heavy mass on her nose. This mass had been present since birth and had gradually increased during puberty without pain or itching. However, it was large enough to be seen by the patient during normal gaze. The patient complained her nose was too heavy and it usually caused a headache. So, she requested to remove it strongly.

On physical examination, the 6.9×6.0×3.5 cm sized dark-brownish mass vertically involved tissue from the nasal tip to the nasal root and the right nasal wall and left cheek (Fig. 1). Punch biopsy was performed pre-

Received: 2021. 2. 22, Revised: 2021. 3. 2, Accepted: 2021. 3. 4
Corresponding author: Gyu Yong Jung, Department of Plastic and Reconstructive Surgery, Dongguk University College of Medicine, 87 Dongdae-ro, Gyeongju 38067, Korea
Tel: 82-54-770-1460, Fax: 82-54-770-1469
E-mail: jnjbeauty@naver.com

Fig. 1. Photograph showing a 6.9×6.0×3.5 cm sized dark-brownish mass, which had been present since birth and grown since puberty, on the patient's nose.

Fig. 2. A forehead flap was performed for nasal reconstruction after mass removal. The flap contained scalp tissue with hair.

operatively and congenital intradermal nevocellular nevus was diagnosed. Preoperative Doppler ultrasonography was used to identify the left supratrochlear artery of the forehead flap for the expected skin defect. Surgery was performed under general anesthesia. During surgery, we removed all of the dark-brownish mass and confirmed that it did not involve nasal mucosa. Next, the size of the skin defect to be covered was measured, and the flap was designed considering the previously identified vascular pedicle. Inevitably, a very large flap with hair was needed to fill the defect. Considering cosmetic out-

Fig. 3. One week after initial surgery, flap blood circulation appeared satisfactory, but the use of scalp tissue resulted in hair on the nasal tip.

Fig. 4. Alexandrite laser hair removal was performed, but a third-degree burn was made on the nasal tip.

come, the flap was made as thin as possible to avoid a bulky nose. The resulting flap was rotated to the medial side and the skin defect was covered (Fig. 2). At the same time, the donor site was repaired by primary closure. Flap viability, blood circulation, and warmth were closely monitored every 2 hours for 48 hours after surgery. During this period there was no evidence of poor blood circulation.

Seven days after the operation, all sutures were removed and the patient felt tingling and numbness on her forehead and scalp (Fig. 3). She was satisfied with the outcome of the operation, but complained strongly about

Fig. 5. After 8 weeks of conservative treatment, the appearance of the nose was as good as that of a normal flap with respect to thickness, texture, color, and sebaceous content.

the hair at the tip of her nose. She requested to remove it as soon as possible. Accordingly, we decided to remove the hair sooner than planned. Epilation was attempted at two weeks postoperatively using an Apogee 6200 Alexandrite LASER (Cynosure, Inc., USA) (40 msec, 18 J) equipped with a Zimmer elektromedizin cooling system.

However, the procedure resulted in a third-degree burn on her nose. A 1.2×1.3 cm size burn wound with eschar was observed (Fig. 4). The patient underwent a second operation three weeks after the first surgery for flap division and inset. The burn on the nasal tip was treated conservatively for eight weeks (Fig. 5). At her most recent follow-up (13 months after the first operation), the patient did not receive any additional hair removal treatment as most of the hair had been removed, and the appearance of her nose was as good as expected of a normal flap with respect to thickness, texture, color, and sebaceous content.

DISCUSSION

Nose is a complex three-dimensional structure and a major aesthetic focus, and thus, nasal reconstruction is the most complex type of facial reconstruction. Several methods are available for treating skin defects in nasal area caused by tumor resection or trauma [2]. The paramedian

forehead flap is the best option for covering large skin defects [3], whereas the forehead flap is an axial-pattern flap with major blood vessels, and branching of these vessels is responsible for blood supply. In general, the supratrochlear or supraorbital artery can be used; in our patient, the left supratrochlear artery was used, based on considerations of flap rotation and location of the lesion. The mass was very large, but fortunately there was no invasion of nasal mucosa or alae. After considering major blood vessels and cosmetic outcomes, we chose to design a vertical flap including scalp rather than an oblique flap without scalp [3]. As a result, scalp tissue with hair was moved to tip of the reconstructed nose. Efforts were made to make the flap as thin as possible to improve cosmetic results, considering flap survival. Although hair follicles of the scalp are deeper than those of other skins, hair follicles are located in the dermis, and thus, their removal risks injuring subdermal plexus. So, we initially planned to use an epilation laser after all procedures were finished. However, the patient complained strongly about the hair at tip of her nose and wanted to remove it promptly. Blood circulation of the flap was the most important concern in terms of flap survival and wound recovery. At one week after initial surgery, the flap was pinkish and its capillary refill time was normal. In the case of full thickness skin graft without axial patterned blood vessels, blood vessels are produced 8 days after transplantation [4]. In our patient, the supratrochlear artery was preserved. So, blood circulation was not considered to be an issue of concern. Hair removal was performed using an Alexandrite laser two weeks after the initial surgery during an outpatient visit. Laser power was adjusted with consideration of the relatively deep hair follicles of the scalp and the hair thickness. Unfortunately, a third-degree burn was occurred on the nasal tip although the same power was applied. The about 1.2×1.3 cm size burn was treated conservatively and healed slowly over about 8 weeks.

We make the following comments regarding the cause of the burn. First of all, clinician's judgment is critical because no research has been conducted on the optimal timing of hair removal after reconstruction. In general, we believe that two weeks is inadequate for flap evaluation. In our case careful monitoring produced no evidence of

poor blood circulation [5]. Because the flap had an axial-patterned blood vessel supply, we considered two weeks would be enough to ensure flap survival. However, when we compared burned and unburned areas, blood vessels that reached the nasal tip after the flap had settled were furthest from the new blood supply and the supratrochlear artery. Furthermore, subsequently it was also apparent that the blood circulation was not enough to promote normal wound healing as it took 8 weeks to heal the 1.2×1.3 cm sized third-degree burn. We used conventional setting for the procedure in consideration of the patient's hair characteristics. This is a setting that does not cause burns a month after the surgery in the previous study [6]. Although the monitoring of the forehead flap was properly performed, two weeks was not enough to endure laser epilation, which made the treatment period longer.

Hair growth on a nose reconstructed using a forehead flap no longer presents a therapeutic or cosmetic challenge. However, in many cases, patients want to remove the hair as soon as possible, whereas operators want to remove the hair as late as possible to ensure flap survival. Further study is needed to determine the appropriate time to perform laser hair removal on reconstructed noses and to provide guidance on the restoration of proper blood circulation for wound healing after hair removal. If hair removal will be performed early after forehead flap surgery, even if blood circulation looks good, it is not recommended to use Alexandrite laser two weeks after the surgery, and it is recommended to wait at least four weeks before performing it. We hope this case study will be helpful by those contemplating hair removal on a flap tip.

CONFLICTS OF INTEREST

None declared.

SOURCE OF FUNDING

None declared.

ACKNOWLEDGEMENT

No benefits in any form have been received or will be received from any commercial party related directly or indirectly to the subject of this article.

REFERENCES

1. Gan SD, Graber EM. Laser hair removal: a review. *Dermatol Surg* 2013;39:823-38.
2. Guo L, Pribaz JR, Pribaz JJ. Nasal reconstruction with local flaps: a simple algorithm for management of small defects. *Plast Reconstr Surg* 2008;122:130e-139e.
3. Correa BJ, Weathers WM, Wolfswinkel EM, Thornton JF. The forehead flap: the gold standard of nasal soft tissue reconstruction. *Semin Plast Surg* 2013;27:96-103.
4. Zarem HA, Zweifach BW, McGehee JM. Development of microcirculation in full thickness autogenous skin grafts in mice. *Am J Physiol* 1967;212:1081-5.
5. Cervenka B, Bewley AF. Free flap monitoring: a review of the recent literature. *Curr Opin Otolaryngol Head Neck Surg* 2015;23:393-8.
6. Yen CI, Chang CJ, Chang CS, Chen HC, Chien MH, Hsiao YC. Laser hair removal following forehead flap for nasal reconstruction. *Lasers Med Sci* 2020;35:1549-54.